



**SQUARE**

$P = 4s$   
 $A = s^2$




**RECTANGLE**

$P = 2a + 2b$   
 $A = ab$




**CIRCLE**

$P = 2\pi r$   
 $A = \pi r^2$




**TRIANGLE**

$P = a + b + c$   
 $A = \frac{1}{2}bh$




**PARALLELOGRAM**

$P = 2a + 2b$   
 $A = bh$




**CIRCULAR SECTOR**

$L = \pi r \frac{\theta}{180^\circ}$   
 $A = \pi r^2 \frac{\theta}{360^\circ}$




**PYTHAGOREAN THEOREM**

$a^2 + b^2 = c^2$   
 $c = \sqrt{a^2 + b^2}$




**CIRCULAR RING**

$A = \pi(R^2 - r^2)$




**SPHERE**

$S = 4\pi r^2$   
 $V = \frac{4\pi r^3}{3}$




**TRAPEZOID**

$P = a + b + c + d$   
 $A = h \frac{a+b}{2}$




**RECTANGULAR BOX**

$A = 2ab + 2ac + 2bc$   
 $V = abc$




**RIGHT CIRCULAR CONE**

$A = \pi r^2 + \pi r s$   
 $s = \sqrt{r^2 + h^2}$   
 $V = \frac{1}{3} \pi r^2 h$




$V = \frac{1}{3} B h$   
 $L.A. = \frac{1}{2} l p$   
 $S.A. = L.A. + B$




**CUBE**

$A = 6l^2$   
 $V = l^3$




**CYLINDER**

$A = 2\pi r(r + h)$   
 $V = \pi r^2 h$



**FRUSTUM OF A CONE**

$V = \frac{1}{3} \pi h (r^2 + rR + R^2) = \frac{1}{3} h (S^2 + s^2 + \sqrt{Ss})$



created by m.d khosrotash

**KHOSROTASH MATH**

<http://khosrotash.ir/>  
[info@khosrotash.com](mailto:info@khosrotash.com)

**REGULAR POLYGON OF n SIDES**

central angle:  $\frac{2\pi}{n}$   
 $A = \frac{1}{4} n a^2 \cot \frac{\pi}{n}$   
 $P = n a$

